



Medium chain triglycerides mix (MCT)
mostly obtained from coconut.

» INDICATIONS

Indicated for supportive therapy in cases of:

- Short bowel syndrome
- Pancreatopathy
- Cystic fibrosis
- Hypercatabolic states (malnutrition, burns, sepsis)
- Lymphedema
 - improves lymph composition
 - does not overload the lymphatic system
- Oxidative disorders of long-chain fatty acids: LCHAD, VLCAD
- MCT-based ketogenic diet, used in the treatment of drug-resistant epilepsy, GLUT-1 deficiency, PDHD deficiency
- Chylothorax
- Other diseases that alter the functionality of chyliferous vessels

» DOSAGE AND INSTRUCTIONS FOR USE

Following medical instructions, taking into account age, weight, and clinical conditions, the caloric portion indicatively provided by MCT should represent around 55% of the fat-derived caloric share. Normally in split doses of 15-20 ml (3-4 teaspoons). MCT Oil can be consumed both raw and heated ($T^{\circ} < 150-160^{\circ}C$, to prevent thermal degradation).

» STORAGE CONDITIONS

Store in a cool and dry place. The expiry date refers to the product in an unopened, correctly stored package.

» IMPORTANT WARNINGS

MCT oil must be taken following medical instruction and monitoring. MCT oil must not be used as one's only source of food. Keep out of the reach of children. Store in a cool and dry place. The expiry date refers to the product in an unopened, correctly stored package.

NUTRITIONAL INFORMATION

food for special medical purposes

	per 100 g
Energy kJ/kcal	3470/830
Fats	100 g
of which saturated fatty acids	100 g
of which caprylic acid (C8:0)	55 g
of which capric acid (C10:0)	43 g
of which caproic acid (C6:0)	0.5 g
of which lauric acid (C12:0)	0.9 g
of which myristic acid (C14:0)	0.6 g
Carbohydrates	0 g
of which sugars	0 g
Protein	0 g
Salt	0 g
Dietary fiber	0 g

DENSITY

1 ml of Chiloil corresponds to 0.95 g