



Hydrothermally treated oat flakes to increase endogenous production of antisecretory protein (antisecretory factor) and increase the balance of cellular liquids.

- » **INDICATIONS** Dietary treatment in children over 3 years of age and in adults in the case of hydrops or Ménière's disease.

- » **DOSAGE AND INSTRUCTIONS FOR USE** It is recommended to take 1 g/kg/ day in 2-3 daily doses, following the physician's prescription.
SPC-Flakes can be taken as is, or mixed in water or yoghurt. SPC-Flakes can also be cooked and eaten as broth or vegetable or legume soup, or added to pasta or any other food. It can also be used in the preparation of oven-baked products such as bread, focaccias, or cookies. The product as is cannot be toasted.

- » **STORAGE CONDITIONS** Store in a cool and dry place.
The expiry date refers to the product correctly stored in a cool and dry place.

- » **IMPORTANT WARNINGS** Use under medical supervision.
SPC-flakes must not be used as one's only food source.
Contains gluten.
SPC-flakes are GMO-free.
Keep out of the reach of children.

NUTRITIONAL INFORMATION
food for special medical purposes

| | per 100 g of flakes |
|--------------------------------|---------------------|
| Energy kj/kcal | 1580/375 |
| Fats | 10 g |
| of which saturated fatty acids | 1.5 g |
| Carbohydrates | 59 g |
| of which sugars | 3 g |
| Fiber | 7.5 g |
| Protein | 12 g |
| Salt * | 35 mg |
| VITAMINS | |
| Biotin | 16 µg |
| Folic acid | 166 µg |
| Niacin | 2.7 mg |
| Pantothenic acid | 1.7 mg |
| Riboflavin | 1.3 mg |
| Thiamine | 0.6 mg |
| Vitamin B6 | 0.1 mg |
| Vitamin B12 | 0.3 µg |
| Vitamin E | 0.7 mg |
| Vitamin K | 1.6 µg |
| MINERALS | |
| Calcium | 88 mg |
| Chloride | 86 mg |
| Chromium | 14 µg |
| Iron | 4.4 mg |
| Phosphorus | 400 mg |
| Iodine | 0.5 µg |
| Magnesium | 130 mg |
| Manganese | 3.1 mg |
| Molybdenum | 130 µg |
| Potassium | 310 mg |
| Selenium | 8 µg |
| Sodium | 14 mg |
| Zinc | 2.4 mg |

* The salt content is due exclusively to the sodium in the product.